



MULTI LIGAND CONTROL-TRI LEVEL
LOT# MLAC1D3

PRODUCT CODE: ML-300B
EXP: 2026-04-18

INTENDED USE

The Multi-ligand Controls are intended for use as an assayed quality control material to monitor the consistency of performance of laboratory test procedures associated with determination and monitoring of the clinical status. This product is a human-serum based, lyophilized control, stabilized with preservatives and can be used with all ELISA and CLIA methods.

SUMMARY AND EXPLANATION

The use of quality control material to assist in the assessment of precision in the clinical laboratory is an integral part of laboratory practices. Controls that contain varied levels of analytes are necessary to insure precision and accuracy in immunoassay systems.

REAGENTS

Monobind's Multi-ligand Controls are intended to be used in the exact manner as patient samples. The control is packaged as 6 vials of 3.0 ml, dried. The analyte activities are adjusted to concentrations in the low, middle and high range in order to monitor the efficacy of the procedure in use.

INSTRUCTIONS FOR USE

- 1) Bring the vials to room temperature before use.
- 2) Carefully unscrew and remove cap.
- 3) Add three (3) ml of distilled or deionized water to each vial. Close the cap tightly and let the contents mix thoroughly for 30 minutes
- 4) Aliquot the materials in 0.5 ml aliquots in cryo vials and store at -20°C.

STORAGE, STABILITY AND DISPOSAL

This product will be stable until the expiration date when stored unopened at 2 to 8°C. Once the control is reconstituted, all analytes will be stable for 7 days when stored tightly capped at 2 to 8°C with the following exceptions: 1) **C-Peptide, f-PSA, and PRL** should be assayed immediately after reconstitution, and 2) **Folate, Insulin, and PRL-Seq** will be stable for 1 day. To avoid contamination, it is recommended labs aliquot required quantities into vials before each use.

After reconstituting, controls should be tightly capped and returned to refrigerator 2 to 8° C as soon as practical after use. (Long term room temperature storage is not supported.) After reconstituting, controls should be tightly capped and frozen within 2-hours. **Once thawed, do not refreeze the control; discard remaining material.** It is recommended that customers aliquot control into separate containers before freezing to allow for use on different days. Outdated material should be discarded as a biohazardous component.

| STORAGE | STABILITY | TEMPERATURE |
|-----------------------|------------------|-------------|
| Lyophilized, Unopened | Three (3) years | < 8°C |
| Reconstituted, Opened | Seven (7) days | 2 - 8°C |
| Reconstituted, Opened | Ninety (90) days | < -10°C |

EXPECTED RANGE OF VALUES

The mean values printed in this insert were derived from replicate analyses and are specific for this lot of product. The tests listed were performed by Monobind QA using representative lots of this product, as well as those of Monobind's AccuBind® ELISA and AccuLite® CLIA reagents.

Individual laboratory means should fall within the corresponding acceptable range; however laboratory means may vary from the listed values during the life of this control. Therefore, each laboratory should establish its own means and acceptable ranges for the product used, using Monobind's assignment only as guide. A trend log should be maintained for batch to batch consistency of the test. Variations over time and between laboratories may be caused by a) differences in laboratory personnel, b) improper technique, c) instrumentation and reagents, d) improper dilutions from the stated manufacturer's procedure, and/ or e) modifications in the manufacturer's test procedure.

Refer to <http://www.monobind.com/site/qc-documents.html> for any updated insert information.

WARNING AND PRECAUTIONS FOR IN VITRO DIAGNOSTIC USE

All products that contain human serum have been found to be non-reactive for HIV 1&2, HIV-Ag, HBsAg, HCV and RPR by FDA required tests. Since no known test can offer complete assurance that infectious agents are absent, all human serum products should be handled as potentially hazardous and capable of transmitting disease. Good laboratory procedures for handling blood products can be found in the Center for Disease Control / National Institute of Health, "Biosafety in Microbiological and Biomedical Laboratories," 2nd Edition, 1988, HHS Publication No. (CDC) 88-8395.

Revision: 1 Date: 2023-06-15 Product Code: ML-300B

For Orders and Inquires, please contact
Monobind Inc.
 100 North Pointe Drive
 Lake Forest, CA 92630 USA

Tel: +1 949.951.2665 Mail: info@monobind.com
 Fax: +1 949.951.3539 Fax: www.monobind.com



CEPartner4U, Eindhoven 13
 3951 DBMaarn, The Netherlands
 www.cepartner4u.eu

Please visit our website to learn more about our products and services.

Glossary of Symbols (EN 960/ISO 15223)



REF: Catalogue Number
 LOT: Batch Code
 EC: Authorized Rep in European Country
 REP: European Conformity



| DOCUMENT HISTORY | | | |
|---------------------------------|-------------|---------------------------------|----------|
| PREPARED BY: <i>[Signature]</i> | DEPT: QC | VERIFIED BY: <i>[Signature]</i> | DEPT: QA |
| APPROVED BY: <i>[Signature]</i> | DEPT: Admin | EFFECTIVE DATE: 2023-06-15 | |
| REVISION: 1 | | DCO: 1624 | |

| EXPECTED RANGE OF VALUES FOR MULTI-LIGAND CONTROL - TRI LEVEL | | | | |
|---|-----------------|-----------------|------------------|-------------------|
| MASTER LOT: MLAC1D3 | | | | |
| Analyte | A | B | C | Method |
| | Range | Range | Range | |
| Allergy | | | | |
| IgE in IU/ml | 99.49 ± 32.83 | 23.24 ± 7.67 | 158.73 ± 52.38 | MB ACCUBIND ELISA |
| | 93.38 ± 30.81 | 20.30 ± 6.70 | 159.85 ± 52.75 | MB ACCULITE CLIA |
| Anemia | | | | |
| Ferritin in ng/ml | 27.79 ± 9.17 | 81.75 ± 26.98 | 333.07 ± 109.91 | MB ACCUBIND ELISA |
| | 24.94 ± 8.23 | 82.13 ± 27.10 | 364.67 ± 120.34 | MB ACCULITE CLIA |
| Folate in ng/ml | 1.85 ± 0.61 | 7.90 ± 2.61 | 12.84 ± 4.24 | MB ACCUBIND ELISA |
| | 2.45 ± 0.81 | 8.58 ± 2.83 | 13.53 ± 4.46 | MB ACCULITE CLIA |
| Vitamin B12 in pg/ml | 286.89 ± 94.67 | 414.64 ± 136.83 | 1032.44 ± 340.70 | MB ACCUBIND ELISA |
| | 330.71 ± 109.13 | 428.07 ± 141.26 | 1021.96 ± 337.25 | MB ACCULITE CLIA |
| Anemia VAST | | | | |
| (Folate) in ng/ml | 2.71 ± 0.89 | 7.83 ± 2.58 | 12.49 ± 4.12 | MB ACCUBIND ELISA |
| | 2.78 ± 0.92 | 7.76 ± 2.56 | 11.62 ± 3.84 | MB ACCULITE CLIA |
| (Vitamin B12) in pg/ml | 366.57 ± 120.97 | 466.13 ± 153.82 | 978.22 ± 322.81 | MB ACCUBIND ELISA |
| | 346.20 ± 114.25 | 469.55 ± 154.95 | 919.75 ± 303.52 | MB ACCULITE CLIA |
| Bone Metabolism | | | | |
| Vit D Direct in ng/ml | 28.57 ± 9.43 | 46.66 ± 15.40 | 88.77 ± 29.29 | MB ACCUBIND ELISA |
| | 31.25 ± 10.31 | 47.51 ± 15.68 | 141.73 ± 46.77 | MB ACCULITE CLIA |
| Cancer Markers | | | | |
| AFP in ng/ml | 20.70 ± 6.83 | 86.38 ± 28.51 | 190.16 ± 62.75 | MB ACCUBIND ELISA |
| | 20.22 ± 6.67 | 91.03 ± 30.04 | 195.92 ± 64.65 | MB ACCULITE CLIA |
| CEA in ng/ml | 4.02 ± 1.33 | 19.12 ± 6.31 | 45.45 ± 15 | MB ACCUBIND ELISA |
| | 3.98 ± 1.31 | 18.12 ± 5.98 | 48.52 ± 16.01 | MB ACCULITE CLIA |
| CEA Next Generation in ng/ml | 4.28 ± 1.41 | 24.29 ± 8.02 | 73.41 ± 24.23 | MB ACCUBIND ELISA |
| | 3.85 ± 1.27 | 22.94 ± 7.57 | 66.64 ± 21.99 | MB ACCULITE CLIA |
| tPSA in ng/ml | 0.77 ± 0.25 | 3.03 ± 1 | > 10 | MB ACCUBIND ELISA |
| | 0.79 ± 0.26 | 3.40 ± 1.12 | > 10 | MB ACCULITE CLIA |
| tPSA-XS in ng/ml | 1.10 ± 0.36 | 3.63 ± 1.20 | 23.03 ± 7.60 | MB ACCUBIND ELISA |
| | 1 ± 0.33 | 3.59 ± 1.18 | 22.99 ± 7.59 | MB ACCULITE CLIA |
| tPSA in ng/ml | 1.35 ± 0.44 | 4.38 ± 1.44 | 25.63 ± 8.46 | MB ACCUBIND ELISA |
| | 1.13 ± 0.37 | 4.07 ± 1.34 | 24.77 ± 8.18 | MB ACCULITE CLIA |
| Cancer Markers VAST | | | | |
| (CEA) in ng/ml | 3.70 ± 1.22 | 18.45 ± 6.09 | 45.61 ± 15.05 | MB ACCUBIND ELISA |
| | 3.33 ± 1.10 | 16.74 ± 5.53 | 46.28 ± 15.27 | MB ACCULITE CLIA |
| (AFP) in ng/ml | 20.81 ± 6.87 | 92.04 ± 30.37 | 189.19 ± 62.43 | MB ACCUBIND ELISA |
| | 19.70 ± 6.50 | 82.27 ± 27.15 | 184.37 ± 60.84 | MB ACCULITE CLIA |
| (tPSA) in ng/ml | 1.23 ± 0.40 | 4.29 ± 1.42 | 30.77 ± 10.16 | MB ACCUBIND ELISA |
| | 1.08 ± 0.36 | 4.24 ± 1.40 | 29.32 ± 9.68 | MB ACCULITE CLIA |
| Cardiac Markers | | | | |
| Dig in ng/ml | 0.36 ± 0.12 | 1.69 ± 0.56 | 2.68 ± 0.88 | MB ACCUBIND ELISA |
| | 0.46 ± 0.15 | 1.61 ± 0.53 | 2.84 ± 0.94 | MB ACCULITE CLIA |
| Diabetes | | | | |
| C-Peptide in ng/ml | 0.48 ± 0.16 | 2.38 ± 0.79 | 4.56 ± 1.50 | MB ACCUBIND ELISA |
| | 0.44 ± 0.15 | 2.29 ± 0.75 | 4.19 ± 1.38 | MB ACCULITE CLIA |
| Insulin in µIU/ml | 28.97 ± 9.56 | 82.75 ± 27.31 | 169.78 ± 56.03 | MB ACCUBIND ELISA |
| | 26.90 ± 8.88 | 84.63 ± 27.93 | 162.90 ± 53.76 | MB ACCULITE CLIA |
| Rapid Insulin in µIU/ml | 28.02 ± 9.25 | 81.63 ± 26.94 | 159.35 ± 52.59 | MB ACCUBIND ELISA |
| | | | | |
| Fertility | | | | |
| FSH in mIU/ml | 8.64 ± 2.85 | 24.11 ± 7.96 | 42.71 ± 14.09 | MB ACCUBIND ELISA |
| | 7.92 ± 2.61 | 23.58 ± 7.78 | 41.52 ± 13.70 | MB ACCULITE CLIA |
| hCG in mIU/ml | 4.43 ± 1.46 | 24.10 ± 7.95 | 146.28 ± 48.27 | MB ACCUBIND ELISA |
| | 4.35 ± 1.74 | 22.95 ± 7.57 | 151.35 ± 49.95 | MB ACCULITE CLIA |
| hCG-XR in mIU/ml | 4.18 ± 1.38 | 28.67 ± 9.46 | 143.88 ± 47.48 | MB ACCUBIND ELISA |
| | 3.56 ± 1.17 | 28.58 ± 9.43 | 155.85 ± 51.43 | MB ACCULITE CLIA |
| LH in mIU/ml | 3.88 ± 1.28 | 22.25 ± 7.34 | 53.53 ± 17.67 | MB ACCUBIND ELISA |
| | 3.38 ± 1.12 | 20.07 ± 6.62 | 53.08 ± 17.72 | MB ACCULITE CLIA |
| PRL in ng/ml | 5.08 ± 1.68 | 24.36 ± 8.04 | 38.50 ± 12.70 | MB ACCUBIND ELISA |
| | 4.78 ± 1.58 | 21.14 ± 6.98 | 36.80 ± 12.14 | MB ACCULITE CLIA |
| PRL-seq in ng/ml | 4.24 ± 1.41 | 20.16 ± 6.65 | 35.96 ± 11.87 | MB ACCUBIND ELISA |
| | 4.14 ± 1.37 | 18.63 ± 6.15 | 36.61 ± 12.08 | MB ACCULITE CLIA |
| Rapid HCG in mIU/ml | 4.70 ± 1.55 | 27.56 ± 9.10 | 188.52 ± 62.21 | MB ACCUBIND ELISA |
| | | | | |
| Fertility VAST | | | | |
| (FSH) in mIU/ml | 6.84 ± 2.26 | 18.59 ± 6.13 | 32.98 ± 10.88 | MB ACCUBIND ELISA |
| | 6.12 ± 2.02 | 17.63 ± 5.82 | 37 ± 12.21 | MB ACCULITE CLIA |
| (LH) in mIU/ml | 4.29 ± 1.42 | 22.36 ± 7.38 | 50.17 ± 16.56 | MB ACCUBIND ELISA |
| | 3.74 ± 1.23 | 20.37 ± 6.72 | 43.9 ± 14.49 | MB ACCULITE CLIA |
| (hCG) in mIU/ml | 4.88 ± 1.61 | 24.08 ± 7.95 | 144.63 ± 47.73 | MB ACCUBIND ELISA |
| | 5.92 ± 1.95 | 26.53 ± 8.76 | 149.81 ± 49.44 | MB ACCULITE CLIA |
| Triple Screen VAST | | | | |
| (AFP) in ng/ml | 21.47 ± 7.09 | 103.20 ± 34.05 | 188.63 ± 62.25 | MB ACCUBIND ELISA |
| | 19.13 ± 6.31 | 100.30 ± 33.10 | 203.38 ± 67.11 | MB ACCULITE CLIA |
| (uE3) in ng/ml | 1.11 ± 0.37 | 3.32 ± 1.10 | 5.99 ± 1.98 | MB ACCUBIND ELISA |
| | 1.10 ± 0.36 | 2.72 ± 0.90 | 5.40 ± 1.78 | MB ACCULITE CLIA |
| (hCG) in mIU/ml | 4.29 ± 1.41 | 23.43 ± 7.73 | 149.04 ± 49.18 | MB ACCUBIND ELISA |
| | 4.78 ± 1.58 | 21.30 ± 7.03 | 174.85 ± 57.70 | MB ACCULITE CLIA |
| Growth Deficiency | | | | |
| hGH in µIU/ml | 5.29 ± 1.75 | 32.33 ± 10.67 | 67.95 ± 22.42 | MB ACCUBIND ELISA |
| | 5.01 ± 1.65 | 32.03 ± 10.57 | 68 ± 23 | MB ACCULITE CLIA |
| Steroids | | | | |
| Aldosterone in ng/ml | 51.72 ± 17.43 | 471.16 ± 155.48 | 1195.18 ± 394.41 | MB ACCUBIND ELISA |
| | 60.35 ± 19.92 | 447.70 ± 147.74 | 1167.75 ± 385.36 | MB ACCULITE CLIA |
| ANST in ng/ml | 1 ± 0.33 | 1.52 ± 0.50 | 10.60 ± 3.50 | MB ACCUBIND ELISA |
| | 0.89 ± 0.29 | 1.36 ± 0.45 | 11.45 ± 3.78 | MB ACCULITE CLIA |
| Cortisol in µg/dl | 2.43 ± 0.80 | 13.98 ± 4.61 | 30.98 ± 11.40 | MB ACCUBIND ELISA |
| | 3.02 ± 1 | 14.91 ± 4.92 | 33.37 ± 11.01 | MB ACCULITE CLIA |
| DHEA-S in µg/ml | 0.37 ± 0.12 | 1.64 ± 0.54 | 4.40 ± 1.45 | MB ACCUBIND ELISA |
| | 0.40 ± 0.17 | 1.51 ± 0.50 | 3.99 ± 1.32 | MB ACCULITE CLIA |
| DHEA in ng/ml | 0.89 ± 0.30 | 2.94 ± 0.97 | 12.42 ± 4.10 | MB ACCUBIND ELISA |
| | 1.02 ± 0.34 | 3.34 ± 1.10 | 14.14 ± 4.67 | MB ACCULITE CLIA |
| E1 in ng/ml | 32 ± 13.02 | 149.61 ± 49.37 | 365.28 ± 120.54 | MB ACCUBIND ELISA |
| | 36.26 ± 11.96 | 180.72 ± 59.64 | 295.83 ± 97.62 | MB ACCULITE CLIA |
| E2 in pg/ml | 35.85 ± 11.83 | 189 ± 62.37 | 283.05 ± 93.41 | MB ACCUBIND ELISA |
| | 1.04 ± 0.41 | 2.43 ± 0.80 | 5.14 ± 1.70 | MB ACCUBIND ELISA |
| uE3 in ng/ml | 1.19 ± 0.39 | 2.51 ± 0.83 | 4.97 ± 1.64 | MB ACCULITE CLIA |
| | 0.97 ± 0.33 | 2.20 ± 0.73 | 25.05 ± 8.27 | MB ACCUBIND ELISA |
| Progesterone in ng/ml | 1.01 ± 0.33 | 7.10 ± 2.34 | 25.39 ± 8.38 | MB ACCULITE CLIA |
| | 0.62 ± 0.20 | 2.01 ± 0.66 | 5.67 ± 1.87 | MB ACCUBIND ELISA |
| 17-OHP in ng/ml | 0.71 ± 0.24 | 2.07 ± 0.68 | 5.71 ± 1.89 | MB ACCULITE CLIA |
| | 0.36 ± 0.12 | 1.13 ± 0.37 | 3 ± 0.99 | MB ACCUBIND ELISA |
| 17-OHP-SI in ng/ml | 0.4 ± 0.13 | 1 ± 0.33 | 2.90 ± 0.96 | MB ACCULITE CLIA |
| | 0.28 ± 0.09 | 1.03 ± 0.34 | 6.93 ± 2.29 | MB ACCUBIND ELISA |
| Testosterone in ng/ml | 0.42 ± 0.14 | 0.90 ± 0.30 | 7.93 ± 2.62 | MB ACCULITE CLIA |
| | 1.11 ± 0.37 | 3.46 ± 1.14 | 28.89 ± 9.53 | MB ACCUBIND ELISA |
| Free Testosterone (0-60pg/ml calibration) | 1.21 ± 0.40 | 3.69 ± 1.22 | 31.29 ± 10.32 | MB ACCULITE CLIA |
| | | | | |
| Thyroid | | | | |
| T3 in ng/ml | 0.51 ± 0.17 | 1.15 ± 0.38 | 3.27 ± 1.08 | MB ACCUBIND ELISA |
| | 0.52 ± 0.17 | 1.17 ± 0.39 | 3.17 ± 1.05 | MB ACCULITE CLIA |
| T4 in µg/dl | 2.90 ± 0.96 | 7.53 ± 2.48 | 16.91 ± 5.58 | MB ACCUBIND ELISA |
| | 2.90 ± 0.96 | 8.37 ± 2.76 | 16.42 ± 5.42 | MB ACCULITE CLIA |
| TSH in µIU/ml | 0.97 ± 0.32 | 6.50 ± 2.14 | 34.20 ± 11.29 | MB ACCUBIND ELISA |
| | 0.88 ± 0.29 | 6.15 ± 2.03 | 31.97 ± 10.55 | MB ACCULITE CLIA |
| fT3 in pg/ml | 1.58 ± 0.52 | 3.46 ± 1.14 | 6.58 ± 2.17 | MB ACCUBIND ELISA |
| | 1.62 ± 0.78 | 3.52 ± 1.16 | 6.78 ± 2.24 | MB ACCULITE CLIA |
| fT4 in ng/dl | 0.38 ± 0.12 | 1.76 ± 0.58 | 3.73 ± 1.23 | MB ACCUBIND ELISA |
| | 0.38 ± 0.12 | 1.63 ± 0.54 | 3.23 ± 1.07 | MB ACCULITE CLIA |
| T3-Uptake in %U | 25.49 ± 2.81 | 33.22 ± 2.94 | 46.13 ± 2.95 | MB ACCUBIND ELISA |
| | 26.63 ± 2.37 | 34.60 ± 2.43 | 49 ± 6.93 | MB ACCULITE CLIA |
| Rapid TSH in µIU/ml | 0.87 ± 0.29 | 6.45 ± 2.13 | 34.26 ± 11.31 | MB ACCUBIND ELISA |
| | 0.77 ± 0.25 | 6.18 ± 2.04 | 31.20 ± 10.30 | MB ACCULITE CLIA |
| Thyroid VAST | | | | |
| (TSH) in µIU/ml | 0.98 ± 0.32 | 7.08 ± 2.34 | 38.39 ± 12.67 | MB ACCUBIND ELISA |
| | 0.94 ± 0.34 | 6.98 ± 2.30 | 35.98 ± 11.87 | MB ACCULITE CLIA |
| Strep T3 in ng/ml | 0.56 ± 0.18 | 1.29 ± 0.43 | 2.88 ± 0.95 | MB ACCUBIND ELISA |
| | 0.63 ± 0.22 | 1.24 ± 0.48 | 2.65 ± 0.88 | MB ACCULITE CLIA |
| Strep T4 in µg/dl | 2.90 ± 0.96 | 9.11 ± 3.01 | 13.56 ± 4.47 | MB ACCUBIND ELISA |
| | 3.08 ± 1.02 | 9.32 ± 3.08 | 12.51 ± 4.13 | MB ACCULITE CLIA |
| Free Thyroid VAST | | | | |
| (TSH) in µIU/ml | 0.79 ± 0.26 | 7.37 ± 2.43 | 33.82 ± 11.16 | MB ACCUBIND ELISA |
| | 0.78 ± 0.26 | 7.23 ± 2.39 | 33.38 ± 11.01 | MB ACCULITE CLIA |
| Strep fT3 in pg/ml | 1.43 ± 0.47 | 3.86 ± 1.27 | 8.24 ± 2.72 | MB ACCUBIND ELISA |
| | 1.59 ± 0.52 | 4.12 ± 1.36 | 8.26 ± 2.7 | |